

PATHOLOGICAL CHANGES IN BLOOD CELLS IN HERBICIDE POISONING

Sotimova Shohida Mansur daughter

Kimyo International university in Tashkent

Abstract The aim of this work was to elucidate the structural bases of the reaction of blood system cells in acute and chronic herbicide poisoning and to find spores for their correction with metal-containing biocomplexes. Based on the goal, the following tasks were set: -to study the morphofunctional state of various (copper-containing cobalt) biocomplexes on the morphofunctional state of the cells of the blood system in acute and chronic herbicide poisoning.

Keywords Herbicide Poisoning. Blood System

Relevance The development of industrial and household chemicals, the intensive use of various chemicals in agriculture has created a new problem for physicians and environmentalists. The expansion of the use of pesticides has led to environmental pollution and an increase in the number of people in contact with these drugs. This contributed to the development of various acute and chronic lesions, the manifestation of mutagenic, allergenic and other undesirable pathological effects. One of the most sensitive tissues to toxic effects is hematopoietic tissue, which contains a pool of rapidly proliferating cells. It clearly manifests the negative effects of pesticides, which determine the value of its study in elucidating the mechanisms of action of one or another pesticide. The works devoted to the study of the pathomorphological state of the cells of the blood system in case of pesticide poisoning do not allow revealing the patterns of changes in the blood System. The development of rational methods of prevention and pathogenetic methods for correcting the toxic effects of pesticides is one of the urgent tasks of modern medicine. Unfortunately, so far there are only a few reports on the use of drugs that have an antidote and therapeutic effect in case of poisoning with certain pesticides. In this regard, the most promising are biocomplexes of various microelements with organic compounds, which have a high protective therapeutic effect in case of poisoning with pesticides and ionizing radiation. However, the effect of these biocomplexes on the body, in particular, on the blood system during acute and chronic administration of organophosphorus pesticides, remains not fully understood. The foregoing indicates the relevance and timeliness

of the study of the morphological and functional characteristics of the response of cells in acute and chronic herbicide poisoning and its correction with Biocomplexes.

Purpose and objectives of the study:

The aim of this work was to elucidate the structural bases of the reaction of blood system cells in acute and chronic herbicide poisoning and to find spores for their correction with metal-containing biocomplexes. Based on the goal, the following tasks were set: - to study the morphofunctional state of various (copper-containing cobalt) biocomplexes on the morphofunctional state of the cells of the blood system in acute and chronic herbicide poisoning. Scientific novelty of the work: In the work, for the first time, using complex general morphological, hematological, cytochemical, radioautographic, and electron microscopic methods, the pathomorphological bases of the reaction of blood cells and hematopoiesis to acute and chronic herbicide poisoning were studied. General and specific patterns of pathomorphological changes in the blood system in the dynamics of acute and chronic intoxication with herbicides have been established. It was found that the most characteristic changes in the system blood in case of herbicide poisoning are anemia, leukocytosis with a shift to the left and thrombocytopenia of cellular and subcellular mechanisms, which constitute inhibition of proliferation and differentiation of hematopoietic cells in the bone marrow, increased destructive changes in their subcellular organelles, and as a result, disorganization of metabolic processes in circulating blood cells. For the first time, metal-containing biocomplexes were tested, which have a pronounced protective and therapeutic effect in acute and chronic herbicide poisoning, an appropriate biocomplex was established and recommended for use in practice.

Scientific and practical value of the study:

The obtained data on shifts in blood parameters and cytochemical changes in neutrophilic leukocytes are additional differential diagnostic tests in the examination of persons working with the herbicide. Data on the cellular and sub-cellular bases of the reaction of blood cells in case of herbicide poisoning can be used in the educational process at medical institutes for a more in-depth explanation of the body's reactions to various Toxic effects. The facts established in the work on the pronounced antidote and therapeutic effect of Biocomplexes make it possible to recommend these preparations for wide use in conditions of work with Herbicides.

The main scientific provisions made:

I. Structural and functional basic reactions of blood system cells in acute and chronic herbicide poisoning have common patterns and are characterized by suppression of the processes of proliferation and differentiation of hematopoietic cells. Biocomplexes have a pronounced protective and therapeutic effect in both acute and chronic herbicide poisoning. Our work is experimental and performed on laboratory rats. However, there are relatively few data on the cells of the blood system of laboratory rats in the literature, and in the vast majority of studies, only the Quantitative content of certain cellular forms of hematopoiesis is considered. The main indicators of peripheral blood and bone marrow of rats are given in the works. They are mostly close to each other. Thus, the total content of erythroblast cells in the bone marrow is in the range of 17.65% -23.6%, mitosis of erythroid cells – 0.29 0.75. The percentage of stab neutrophils in the bone marrow is relatively high. In the leukocyte formula of the peripheral blood of rats, lymphocytes predominate. Sometimes reaching more than 70% of all leukocytes. Among granulocytes, mainly segmented neutrophils are found, Thus, rats are characterized by a higher content of stab neutrophils in the bone marrow and the Predominance of lymphocytes in the leukocyte formula. Currently, a unitary scheme of hematopoiesis is generally accepted, reflecting the successive stages of Histogenesis of blood cells. It has been established that all blood cells originate from a single ancestral hematopoietic stem cell, which a well-known histologist once wrote about. But, going into the details of the discussion about the morphology of hematopoietic stem cells, we considered it appropriate to present the basic data on the morphology of the identifiable cells of the blood system. The ultrastructure of morphologically identifiable cells of the human blood system and small-nourishing cells has been studied in some detail by now. However, these studies have been performed primarily in humans, mice, and rabbits, while studies Only a small number of works have been devoted to the Ultrastructural features of rat hematopoietic cells.

The process of maturation of the cells of the erythroid germ is characterized by certain ultrastructural changes both in the nucleus and in the cytoplasm of the cells. The nucleus decreases in size, taking on a more rounded shape, while chromatin compaction occurs and the nucleolus disappears. The size and number Of mitochondria decreases and the electron density of the mitochondrial matrix increases due to the accumulation of granular content. The content of free ribosomes and polysomes gradually decreases, while the amount of amphora material increases. The electron density of the cytoplasm increases due to the Accumulation of hemoglobin. It has been established that the disappearance of the nucleus from normoblasts

occurs by its expulsion. Mature erythrocytes have a high electron density and are surrounded by a membrane about 20 nm thick; cell organelles are absent. In the process of maturation of neutrophils, two stages are morphologically distinguished, characterized by the formation of two structurally and chemically different populations of granules. The first stage of granulogenesis occurs at the pro-myelocyte stage, when primary or azurophilic granules are formed. Numerous electron microscopic studies have shown that the changes that occur in the process of Differentiation of human eosinophilic leukocytes and other small-nourishing leukocytes are in many respects Similar to those of neutrophilic leukocytes. The formation of specific eosinophilic granules is carried out in The components of the Golgi complex. A characteristic feature of small-feeding eosinophilic granules is the Presence of an electron-dense crystalloid in their mainland. Electron-cytochemically and biochemically, acid phosphatase, arylsulfatase, peroxidase, esterase, basic Proteins and phospholipids were detected in eosinophilic granules.

The ultrastructure of basophilic leukocytes has been studied in less detail. According to the characteristics of the nucleus and cytoplasm, with the exception of granules, basophilic leukocytes practically do not differ from neutrophilic and eosinophilic ones. Specific basophilic granules vary in size and structure. They are predominantly round or oval in shape, 0.2-1.62 microns in size, surrounded by a membrane. The functional role of basophilic granulocytes has not yet been fully elucidated. It is assumed that they are involved in allergic reactions of the body and in the metabolism of histamine. A number of works have been devoted to the study of the ultrastructure of monocytopoietic cells. Currently, these cells are considered as precursors of tissue macrophages and are united in the system of mononuclear phagocytes. They are characterized by the Presence of granules with the activity of myeloperoxidase and lysosomal enzymes. In connection with the achievement of immunology, there is now a need for research that would simultaneously make it possible to judge whether lymphocytes belong to 1 or 2-immunity systems and evaluate their structural features. A broad perspective for solving this problem is opening up in connection with the introduction of methods of immune electron microscopy, which combines the achievements of morphology and immunochemistry. Studies of functionally heterogeneous rat lymphocytes using immune electron microscopy are still rare and insufficient for the development of morphological criteria for various cell populations. Thus, the analysis of the literature data allows us to conclude that quite extensive information on the Structural and functional features of the cells of the mammalian blood system has been

accumulated to date. However, it reflects the state structures and functions of cells under physiological conditions and in various diseases of the blood system.

The pathomorphological basis of adaptive rearrangements of blood cells and hematopoietic tissue in acute and chronic poisoning with pesticides, in particular, herbicides, remains practically unexplained. Meanwhile, the disclosure of the structural and functional regularities of the effect of herbicides on the cells of the blood system allows us to consider in depth the mechanism of changes observed in the body and contributes to the choice of the most optimal methods for their correction. In terms Of pharmacological correction of the response of cells and tissues to the action of herbicides, preparations are the most promising. The foregoing as a whole necessitates further research, which provides for the elucidation of the structural and functional features of the reactions of blood cells to the action of a widespread pesticide and the Possibilities of pharmacological correction of these reactions using metal-containing biological complexes.

Materials and Research Methods:

The experiments were carried out in 224 outbred male rats with a body weight of 140-150 g. The animals were kept under standard vivarium conditions. Before the experiments, the animals were quarantined for two Weeks to exclude various diseases. All animals were divided into two main groups. The first group of animals (84 rats) was divided into 4 subgroups and was used to study the state of the reaction of the blood system during acute poisoning with herbicides and in the process of its correction with biocomplexes. Three subgroups of this group received a single dose of herbicide intragastrically at a dose of 150 mg/kg. The fourth subgroup, which received an equal volume of sterile saline, served as the control. Animals of subgroup I were slaughtered under ether anesthesia on days 1.3 and 7 after herbicide poisoning. Animals of subgroup II, starting from the first day after poisoning, received biocomplex No. I once a day subcutaneously at a dose of 10 mg/kg for 7 days. Animals of the III subgroup similarly subcutaneously injected biocomplex No. 6 for 7 days at 10 mg/kg. Both (I and II) subgroups of animals were slaughtered on the 7th day of the experiments simultaneously with the control subgroups of rats in compliance with all Conditions.

Conclusions:

With acute exposure to fosalone at a dose of 1/20 LD 50, against the background of signs of intoxication, severe hypochromic anemia develops leukocytosis with a shift to the left and thrombocytopenia. This is accompanied by inhibition of the activity of myeloperoxidase and

alkaline phosphatase in neutrophilic leukocytes, destructive changes in subcellular organelles, and disruption of the process of granulogenesis in maturing granulocytes. An increase in the degree of ineffective erythropoiesis and intramedullary destruction of erythroid cells, a decrease in their proliferative activity are separate links in the pathogenesis of anemia in acute and chronic intoxication with herbicides. In the pathogenesis of thrombocytopenia in acute and chronic poisoning with phosalone, an important role belongs to the destructive changes in the organelles of megakaryocytes, the reduction of their demarcation membranes, and the slowing down of the processes of platelet lacing. For the first time, a comparative study of the membrane-stabilizing and therapeutic properties of new microelement biocomplexes with vitamin U and amino acids was carried out. It has been established that biocomplexes of cobalt and copper have a rather high membrane-stabilizing and hemostimulating activity in both acute and chronic herbicide intoxication. According to the results of cytochemical, radioautographic and electron microscopic studies, it was found that the biocomplex Co (II) is the most harmless and highly effective through the membrane stabilizing effect, helps to restore damaged subcellular organelles and has a stimulating effect on the proliferative activity of cells of the blood system.

Literature

1. Abakeliya Ts.I., Larionova N.R., Shukiya T.Sh., Tsomaya I.S. Morpho-cytochemical characteristics of bone marrow cells of healthy laboratory rats. In the book: Questions of clinical and experimental radiology. Tbilisi, 1975, p. 141-145.
- 2 Akbarov A.B., Tursunov E.A., Nuritov N.R. Aqua methylmethioninate sulfonium, a disubstituted. copper-(II) glutamate, exhibiting antidote properties when etched with phosalone. A.S. No. 1612544 USSR, chipboard, MKI C 07 I 15/04, A 61 K 31/295-6 p.
3. Akbarov A.B., Aliev U.E., Temirkhodzhaev V.Kh.. Tursunov E.A. and others. Tetrahydro-, bio- methylmethioninatesulfonium chloride, monosubstituted cobalt-(II) glutamate, showing ventilatory and anti-radiation activity. A.S. No. 1426059 USSR, DSN, MKI S 07 G 15/06 A 61 K 31/295-6 p.
4. Akilov A.T. Cytohistochemical analysis of hematopoietic foci and elements of peripheral blood in some representatives of vertebrates in different seasons of the year and under the action of ionizing radiation. - Abstract of the thesis. ... doc. biol.sci. Tashkent, 1972, 40 p.

5. Afanasiev Yu.I. Organs of hematopoiesis and immunological protection. In the book: Histology (under the editorship of Eliseev V.G., Afanasiev Yu.I., Yurina N.A.) M., Medicine, 1983, p.345-Date of Publication:03-11-2022370.

6. Ataev M.A. Influence of the pesticide magnesium chlorate on the coagulating activity of blood: Abstract of the thesis.... C.m.s. – Ashgabat, 1980, 25 p.

7. Barkova E.K., Petrov A.V. Influence of erythropoietin on the ultrastructure of erythroblastic elements of the bone marrow. Bull. Exp. Biology and medicine, 1973, v. 76, 7, p. III-114. .

8 Butenko Z.A., Gluzman D.F., Zak K.P. Cytochemistry and electron microscopy of blood cells and hematopoietic organs. – Kyiv: Science Dumka, 1974, 266 p.